

MECOM Break Apart FISH Probe Kit

Introduction

The MECOM Break Apart FISH Probe Kit is designed to detect rearrangements in the human *MECOM* locus located on chromosome band 3q26.2. In addition to revealing breaks, which can lead to translocation of parts of the gene, inversion, or its fusion to other genes, the probe set can also be used to identify other *MECOM* aberrations such as deletions or amplifications. Rearrangements and abnormal expression of the *MECOM* gene – also known as *EVI1*, *MDS1*, *PRDM3*, *MDS1-EVI1* or *AML1-EVI-1* – have been observed in acute and chronic myelogenous leukemias, Myelodysplastic Syndrome (MDS) and other malignancies.

Intended Use


To detect rearrangements in the human *MECOM* gene located on chromosome band 3q26.2.

Cont.	Color
-------	-------

LSP MECOM 5' FISH Probe
LSP MECOM 3' FISH Probe

CytoGreen
CytoOrange

Probe Design



LSP MECOM 5' FISH Probe covers the 5' (start) portion of the *MECOM* gene and some adjacent genomic sequences. LSP MECOM 3' FISH Probe covers the 3' (end) part as well as sequences downstream of the gene. The two probes are flanking sequences across the *MECOM* gene in which variable breakpoints have been observed.

Not to Scale

Cat. No.	Volume
----------	--------

CT-PAC071-10-GO

10 Tests (100 µL)

Signal Pattern Interpretation

Normal Pattern

2OG*

Abnormal Pattern

Other Patterns

*Overlapping orange and green signals can appear as yellow.

- 1) Wieser R, et al. *Haematologica*. 88(1):25-30 (2003).
- 2) Nonet GH, et al. *Cancer Res*. 2001 Feb 15;61(4):1250-4 (2001).
- 3) Poppe B, et al. *Genes Chromosomes Cancer*. 45(4):349-56 (2006).
- 4) Yin CC, et al. *Cancer*. 106(8):1730-8 (2006).
- 5) Bobadilla D, et al. *Br J Haematol*. 136(6):806-13 (2007).



CytoTest Inc.
9430 Key West Ave., Suite 210
Rockville, MD 20850, USA

* CE IVD only available in certain countries. All other countries are either ASR or RUO. Please contact your local dealer or our headquarters for more information.